NUTRITION

Migraine Headaches and Food: The "Trigger Factor"

Up to 20 percent of the 23 million migraine sufferers in this country experience painful, debilitating migraine headaches that are initiated by eating a particular food or combination of foods. Migraine headaches can affect anyone, but women are three times more likely to suffer from them than men. If you are a sufferer of migraine headaches, you need to see a physician for a proper diagnosis and treatment plan. If you have linked your migraine headaches to "trigger" foods, try to avoid them and inform your doctor.

What may cause a migraine attack?

Even though the actual cause of migraines remains uncertain, various factors can "trigger" the attack. Factors that may "trigger" migraines are certain types of foods, hunger of missed meals, changes in weather, some physical activities, emotions and stress, medication and hormonal changes.

List of foods that may trigger a migraine attack

1 41 1 1	D 1 : 41 1 11
1. Alcohol	Red wine, vermouth, champagne and beer

2. Beverages with caffeine Coffee, tea and soft drinks

3. Dairy products Aged cheeses

4. Breads Sourdough, fresh yeast and homemade breads

5. Vegetables & Fruits Peas, avocados, overripe bananas and some types of

beans (Italian, broad, lentil, fava and soy)

6. Snacks Peanuts and peanut butter

7. Meats Salted and cured meats (ham, corned beef and lunch

meats), dried meats, pickled herring and chicken

livers

8. Soups Canned or from mixes9. Desserts Chocolate-based desserts

Carefully planned menus may reduce the incidence of migraine attacks. Remember it is especially important to see your physician for proper diagnosis and treatment.

For a free copy of "Chart Your Route to Relief" from Glaxo Wellcome write P.O. Box 800, Wendell, NC 27591 or call 1-800-377-0302.



Office of Insurance Services